

# World Journal of *Clinical Cases*

*World J Clin Cases* 2020 October 6; 8(19): 4280-4687



**OPINION REVIEW**

- 4280 Role of monoclonal antibody drugs in the treatment of COVID-19  
*Ucciferri C, Vecchiet J, Falasca K*

**MINIREVIEWS**

- 4286 Review of simulation model for education of point-of-care ultrasound using easy-to-make tools  
*Shin KC, Ha YR, Lee SJ, Ahn JH*
- 4303 Liver injury in COVID-19: A minireview  
*Zhao JN, Fan Y, Wu SD*

**ORIGINAL ARTICLE****Case Control Study**

- 4311 Transanal minimally invasive surgery *vs* endoscopic mucosal resection for rectal benign tumors and rectal carcinoids: A retrospective analysis  
*Shen JM, Zhao JY, Ye T, Gong LF, Wang HP, Chen WJ, Cai YK*
- 4320 Impact of *mTOR* gene polymorphisms and gene-tea interaction on susceptibility to tuberculosis  
*Wang M, Ma SJ, Wu XY, Zhang X, Abesig J, Xiao ZH, Huang X, Yan HP, Wang J, Chen MS, Tan HZ*

**Retrospective Cohort Study**

- 4331 Establishment and validation of a nomogram to predict the risk of ovarian metastasis in gastric cancer: Based on a large cohort  
*Li SQ, Zhang KC, Li JY, Liang WQ, Gao YH, Qiao Z, Xi HQ, Chen L*

**Retrospective Study**

- 4342 Predictive factors for early clinical response in community-onset *Escherichia coli* urinary tract infection and effects of initial antibiotic treatment on early clinical response  
*Kim YJ, Lee JM, Lee JH*
- 4349 Managing acute appendicitis during the COVID-19 pandemic in Jiaying, China  
*Zhou Y, Cen LS*
- 4360 Clinical application of combined detection of SARS-CoV-2-specific antibody and nucleic acid  
*Meng QB, Peng JJ, Wei X, Yang JY, Li PC, Qu ZW, Xiong YF, Wu GJ, Hu ZM, Yu JC, Su W*
- 4370 Prolonged prothrombin time at admission predicts poor clinical outcome in COVID-19 patients  
*Wang L, He WB, Yu XM, Hu DL, Jiang H*

- 4380 Percutaneous radiofrequency ablation is superior to hepatic resection in patients with small hepatocellular carcinoma

*Zhang YH, Su B, Sun P, Li RM, Peng XC, Cai J*

- 4388 Clinical study on the surgical treatment of atypical Lisfranc joint complex injury

*Li X, Jia LS, Li A, Xie X, Cui J, Li GL*

- 4400 Application of medial column classification in treatment of intra-articular calcaneal fractures

*Zheng G, Xia F, Yang S, Cui J*

### Clinical Trials Study

- 4410 Optimal hang time of enteral formula at standard room temperature and high temperature

*Lakananurak N, Nalinthassanai N, Suansawang W, Panarat P*

### META-ANALYSIS

- 4416 Meta-analysis reveals an association between acute pancreatitis and the risk of pancreatic cancer

*Liu J, Wang Y, Yu Y*

### SCIENTOMETRICS

- 4431 Global analysis of daily new COVID-19 cases reveals many static-phase countries including the United States potentially with unstoppable epidemic

*Long C, Fu XM, Fu ZF*

### CASE REPORT

- 4443 Left atrial appendage aneurysm: A case report

*Belov DV, Moskalev VI, Garbuzenko DV, Arefyev NO*

- 4450 Twenty-year survival after iterative surgery for metastatic renal cell carcinoma: A case report and review of literature

*De Raffe E, Mirarchi M, Casadei R, Ricci C, Brunocilla E, Minni F*

- 4466 Primary rhabdomyosarcoma: An extremely rare and aggressive variant of male breast cancer

*Satală CB, Jung I, Bara TJ, Simu P, Simu I, Vlad M, Szodorai R, Gurzu S*

- 4475 Bladder stones in a closed diverticulum caused by *Schistosoma mansoni*: A case report

*Alkhamees MA*

- 4481 Cutaneous ciliated cyst on the anterior neck in young women: A case report

*Kim YH, Lee J*

- 4488 Extremely rare case of successful treatment of metastatic ovarian undifferentiated carcinoma with high-dose combination cytotoxic chemotherapy: A case report

*Kim HB, Lee HJ, Hong R, Park SG*

- 4494** Acute amnesia during pregnancy due to bilateral fornix infarction: A case report  
*Cho MJ, Shin DI, Han MK, Yum KS*
- 4499** Ascaris-mimicking common bile duct stone: A case report  
*Choi SY, Jo HE, Lee YN, Lee JE, Lee MH, Lim S, Yi BH*
- 4505** Eight-year follow-up of locally advanced lymphoepithelioma-like carcinoma at upper urinary tract: A case report  
*Yang CH, Weng WC, Lin YS, Huang LH, Lu CH, Hsu CY, Ou YC, Tung MC*
- 4512** Spontaneous resolution of idiopathic intestinal obstruction after pneumonia: A case report  
*Zhang BQ, Dai XY, Ye QY, Chang L, Wang ZW, Li XQ, Li YN*
- 4521** Successful pregnancy after protective hemodialysis for chronic kidney disease: A case report  
*Wang ML, He YD, Yang HX, Chen Q*
- 4527** Rapid remission of refractory synovitis, acne, pustulosis, hyperostosis, and osteitis syndrome in response to the Janus kinase inhibitor tofacitinib: A case report  
*Li B, Li GW, Xue L, Chen YY*
- 4535** Percutaneous fixation of neonatal humeral physal fracture: A case report and review of the literature  
*Tan W, Wang FH, Yao JH, Wu WP, Li YB, Ji YL, Qian YP*
- 4544** Severe fundus lesions induced by ocular jellyfish stings: A case report  
*Zheng XY, Cheng DJ, Lian LH, Zhang RT, Yu XY*
- 4550** Application of ozonated water for treatment of gastro-thoracic fistula after comprehensive esophageal squamous cell carcinoma therapy: A case report  
*Wu DD, Hao KN, Chen XJ, Li XM, He XF*
- 4558** Germinomas of the basal ganglia and thalamus: Four case reports  
*Huang ZC, Dong Q, Song EP, Chen ZJ, Zhang JH, Hou B, Lu ZQ, Qin F*
- 4565** Gastrointestinal bleeding caused by jejunal angiosarcoma: A case report  
*Hui YY, Zhu LP, Yang B, Zhang ZY, Zhang YJ, Chen X, Wang BM*
- 4572** High expression of squamous cell carcinoma antigen in poorly differentiated adenocarcinoma of the stomach: A case report  
*Wang L, Huang L, Xi L, Zhang SC, Zhang JX*
- 4579** Therapy-related acute promyelocytic leukemia with FMS-like tyrosine kinase 3-internal tandem duplication mutation in solitary bone plasmacytoma: A case report  
*Hong LL, Sheng XF, Zhuang HF*
- 4588** Metastasis of esophageal squamous cell carcinoma to the thyroid gland with widespread nodal involvement: A case report  
*Zhang X, Gu X, Li JG, Hu XJ*

- 4595** Severe hyperlipemia-induced pseudoerythrocytosis - Implication for misdiagnosis and blood transfusion: A case report and literature review  
*Zhao XC, Ju B, Wei N, Ding J, Meng FJ, Zhao HG*
- 4603** Novel brachytherapy drainage tube loaded with double 125I strands for hilar cholangiocarcinoma: A case report  
*Lei QY, Jiao DC, Han XW*
- 4609** Resorption of upwardly displaced lumbar disk herniation after nonsurgical treatment: A case report  
*Wang Y, Liao SC, Dai GG, Jiang L*
- 4615** Primary hepatic myelolipoma: A case report and review of the literature  
*Li KY, Wei AL, Li A*
- 4624** Endoscopic palliative resection of a giant 26-cm esophageal tumor: A case report  
*Li Y, Guo LJ, Ma YC, Ye LS, Hu B*
- 4633** Solitary hepatic lymphangioma mimicking liver malignancy: A case report and literature review  
*Long X, Zhang L, Cheng Q, Chen Q, Chen XP*
- 4644** Intraosseous venous malformation of the maxilla after enucleation of a hemophilic pseudotumor: A case report  
*Cai X, Yu JJ, Tian H, Shan ZF, Liu XY, Jia J*
- 4652** Intravesically instilled gemcitabine-induced lung injury in a patient with invasive urothelial carcinoma: A case report  
*Zhou XM, Wu C, Gu X*
- 4660** Bochdalek hernia masquerading as severe acute pancreatitis during the third trimester of pregnancy: A case report  
*Zou YZ, Yang JP, Zhou XJ, Li K, Li XM, Song CH*
- 4667** Localized primary gastric amyloidosis: Three case reports  
*Liu XM, Di LJ, Zhu JX, Wu XL, Li HP, Wu HC, Tuo BG*
- 4676** Displacement of peritoneal end of a shunt tube to pleural cavity: A case report  
*Liu J, Guo M*
- 4681** Parathyroid adenoma combined with a rib tumor as the primary disease: A case report  
*Han L, Zhu XF*

**ABOUT COVER**

Peer-reviewer of *World Journal of Clinical Cases*, Prof. Adrián Ángel Inchauspe, obtained his MD in 1986 from La Plata National University (Argentina), where he remained as Professor of Surgery. Study abroad, at the Aachen and Tübingen Universities in Germany in 1991, led to his certification in laparoscopic surgery, and at the Louis Pasteur University in Strasbourg France, led to his being awarded the Argentine National Invention Award in 1998 for his graduate work in tele-surgery. He currently serves as teacher in the Argentine Acupuncture Society, as Invited Foreigner Professor at the China National Academy of Sciences and Hainan Medical University, and as editorial member and reviewer for many internationally renowned journals. (L-Editor: Filipodia)

**AIMS AND SCOPE**

The primary aim of *World Journal of Clinical Cases* (*WJCC*, *World J Clin Cases*) is to provide scholars and readers from various fields of clinical medicine with a platform to publish high-quality clinical research articles and communicate their research findings online.

*WJCC* mainly publishes articles reporting research results and findings obtained in the field of clinical medicine and covering a wide range of topics, including case control studies, retrospective cohort studies, retrospective studies, clinical trials studies, observational studies, prospective studies, randomized controlled trials, randomized clinical trials, systematic reviews, meta-analysis, and case reports.

**INDEXING/ABSTRACTING**

The *WJCC* is now indexed in Science Citation Index Expanded (also known as SciSearch®), Journal Citation Reports/Science Edition, PubMed, and PubMed Central. The 2020 Edition of Journal Citation Reports® cites the 2019 impact factor (IF) for *WJCC* as 1.013; IF without journal self cites: 0.991; Ranking: 120 among 165 journals in medicine, general and internal; and Quartile category: Q3.

**RESPONSIBLE EDITORS FOR THIS ISSUE**

Production Editor: Yan-Xia Xing; Production Department Director: Yun-Xiaojuan Wu; Editorial Office Director: Jin-Lai Wang.

**NAME OF JOURNAL**

*World Journal of Clinical Cases*

**ISSN**

ISSN 2307-8960 (online)

**LAUNCH DATE**

April 16, 2013

**FREQUENCY**

Semimonthly

**EDITORS-IN-CHIEF**

Dennis A Bloomfield, Sandro Vento, Bao-Gan Peng

**EDITORIAL BOARD MEMBERS**

<https://www.wjgnet.com/2307-8960/editorialboard.htm>

**PUBLICATION DATE**

October 6, 2020

**COPYRIGHT**

© 2020 Baishideng Publishing Group Inc

**INSTRUCTIONS TO AUTHORS**

<https://www.wjgnet.com/bpg/gerinfo/204>

**GUIDELINES FOR ETHICS DOCUMENTS**

<https://www.wjgnet.com/bpg/GerInfo/287>

**GUIDELINES FOR NON-NATIVE SPEAKERS OF ENGLISH**

<https://www.wjgnet.com/bpg/gerinfo/240>

**PUBLICATION ETHICS**

<https://www.wjgnet.com/bpg/GerInfo/288>

**PUBLICATION MISCONDUCT**

<https://www.wjgnet.com/bpg/gerinfo/208>

**ARTICLE PROCESSING CHARGE**

<https://www.wjgnet.com/bpg/gerinfo/242>

**STEPS FOR SUBMITTING MANUSCRIPTS**

<https://www.wjgnet.com/bpg/GerInfo/239>

**ONLINE SUBMISSION**

<https://www.f6publishing.com>

## Left atrial appendage aneurysm: A case report

Dmitry Vladimirovich Belov, Vladimir Ivanovich Moskalev, Dmitry Victorovich Garbuzenko, Nikolay Olegovich Arefyev

**ORCID number:** Dmitry Vladimirovich Belov [0000-0003-4985-9716](https://orcid.org/0000-0003-4985-9716); Vladimir Ivanovich Moskalev [0000-0001-9759-7146](https://orcid.org/0000-0001-9759-7146); Dmitry Victorovich Garbuzenko [0000-0001-9809-8015](https://orcid.org/0000-0001-9809-8015); Nikolay Olegovich Arefyev [0000-0002-1770-064X](https://orcid.org/0000-0002-1770-064X).

**Author contributions:** Belov DV and Moskalev VI collected the data, reviewed the literature and contributed to manuscript drafting; Garbuzenko DV contributed to manuscript drafting and revised manuscript for important intellectual content; Arefyev NO drafted the manuscript and reviewed the literature; all authors issued final approval for the version to be submitted.

**Informed consent statement:** Informed written consent was obtained from the patient for publication of this report and any accompanying images.

**Conflict-of-interest statement:** The authors declare that they have no conflict of interest.

**CARE Checklist (2016) statement:** The authors have read the CARE Checklist (2016), and the manuscript was prepared and revised according to the CARE Checklist (2016).

**Open-Access:** This article is an open-access article that was selected by an in-house editor and

**Dmitry Vladimirovich Belov**, Department of Hospital Surgery, South Ural State Medical University, Chelyabinsk 454092, Russia

**Dmitry Vladimirovich Belov, Vladimir Ivanovich Moskalev**, Department of Hospital Surgery, Federal Center for Cardiovascular Surgery, Chelyabinsk 454033, Russia

**Dmitry Victorovich Garbuzenko**, Department of Faculty Surgery, South Ural State Medical University, Chelyabinsk 454092, Russia

**Nikolay Olegovich Arefyev**, Department of Pathological Anatomy and Forensic Medicine, South Ural State Medical University, Chelyabinsk 454092, Russia

**Corresponding author:** Nikolay Olegovich Arefyev, MD, Academic Research, Surgeon, Department of Pathological Anatomy and Forensic Medicine, South Ural State Medical University, Vorovskogo 64, Chelyabinsk 454092, Russia. [nikolai.arefyev@gmail.com](mailto:nikolai.arefyev@gmail.com)

### Abstract

#### BACKGROUND

An aneurysm of the left atrial appendage is one of the rare but potentially hazardous heart defects. The risk of lethal complications grows with its size. To date, about 150 cases of this defect have been described in the literature. We present a case of left atrial appendage aneurysm with the deformation of the mitral valve and the left main coronary and circumflex artery, which required mitral valve annuloplasty and bifurcation stenting.

#### CASE SUMMARY

A 58-year-old man presented to our hospital complaining of shortness of breath, general weakness, dizziness during physical exertion, and fatigue. Based on the results of echocardiography, an aneurysm of the left atrium was suspected. A free-breathing real-time cine magnetic resonance imaging with electrocardiograph synchronization confirmed the diagnosis of left atrial appendage aneurysm. The patient underwent an aneurysmectomy *via* a median sternotomy with cardiopulmonary bypass. Intraoperative transesophageal echocardiography revealed relative mitral insufficiency that was corrected with an annuloplasty ring. Intraoperative coronary angiogram showed impaired blood flow in the left main coronary and circumflex artery and 60% stenosis. For this reason, bifurcation stenting was performed. The patient had an uneventful postoperative clinical course and was discharged from the hospital on the 10th day in a satisfactory condition.

fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Manuscript source:** Invited manuscript

**Received:** May 16, 2020

**Peer-review started:** May 16, 2020

**First decision:** June 7, 2020

**Revised:** June 8, 2020

**Accepted:** September 10, 2020

**Article in press:** September 10, 2020

**Published online:** October 6, 2020

**P-Reviewer:** Ho CM

**S-Editor:** Zhang L

**L-Editor:** A

**P-Editor:** Xing YX



## CONCLUSION

Left atrial appendage aneurysm is a rare and dangerous heart pathology that requires surgery to prevent related complications.

**Key Words:** Atrial appendage; Aneurysm; Mitral valve insufficiency; Coronary stenosis; Catheter ablation; Case report

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

**Core Tip:** Left atrial appendage aneurysm is one of the rare, but potentially hazardous heart defects. We present a case of left atrial appendage aneurysm resection and mitral valve annuloplasty. This case highlights the fact that the disease is diagnosed either by chance or when examining the patient as a result of complications. Echocardiography in most cases will make it possible to establish the diagnosis, but magnetic resonance imaging is necessary for a more detailed assessment of changes in the heart. Surgical treatment is indicated for all patients to prevent fatal disorders.

**Citation:** Belov DV, Moskalev VI, Garbuzenko DV, Arefyev NO. Left atrial appendage aneurysm: A case report. *World J Clin Cases* 2020; 8(19): 4443-4449

**URL:** <https://www.wjgnet.com/2307-8960/full/v8/i19/4443.htm>

**DOI:** <https://dx.doi.org/10.12998/wjcc.v8.i19.4443>

## INTRODUCTION

Left atrial appendage aneurysm is one of the rare but potentially hazardous heart defects. The bigger it grows, the higher is the risk of lethal complications<sup>[1]</sup>. The paper presents a case of left atrial appendage aneurysm that required a left atrial appendage resection and mitral valve annuloplasty.

The PubMed database, the Google Scholar retrieval system, and the reference lists from related articles were used to search for relevant publications. Articles corresponding to the aim of the review were selected for 1962 to 2019 using the keywords: "left atrial appendage aneurysm", "left atrial appendage dilatation", "pathogenesis", "diagnosis", "treatment". The management of patients with left atrial appendage aneurysm was the inclusion criterion.

## CASE PRESENTATION

### Chief complaints

A 58-year old man was hospitalized to the Federal Center for Cardiovascular Surgery of the Russian Ministry of Health (Chelyabinsk) complaining of dyspnea, general weakness, dizziness during physical exertion, and fatigue.

### History of present illness

Over the past five years, the patient experienced rare heart palpitations that have occurred and stopped on their own after a few minutes. Six months before hospitalization, shortness of breath, weakness, and post-exertion dizziness appeared. Heart rate increased to 120 per minute. The electrocardiogram taken in an outpatient clinic revealed an irregular form of atypical atrial flutter. Based on the results of echocardiography, an aneurysm of the left atrium was suspected. Warfarin was prescribed. The maintenance dose was 2.5 mg daily.

### History of past illness

The patient had a free previous medical history.

### Physical examination

On the patient's visit to the Center for Cardiovascular Surgery, his vital signs were within normal limits.

### Laboratory examinations

The Quick prothrombin activity was 27.9% (reference values were 70%-120%), INR was 2.52, and the prothrombin time was 45.6 s due to warfarin treatment. Other laboratory parameters of blood and urine analyses, blood biochemistry, and arterial blood gas were within normal limits.

### Imaging examinations

A free-breathing real-time cine magnetic resonance imaging (MRI) with electrocardiograph synchronization did not reveal any defects in the leaves of the pericardium and extrapericardial protrusions of the heart cavities. A giant aneurysm of the left atrial appendage (123 mm × 90 mm × 70 mm in size) had uneven and clear contours, thin walls, and turbulent blood flow in the cavity. The aneurysm had a wide communication with the cavity of the left atrium. The neck of the aneurysm was 51 mm × 55 mm in size. The cavity of the aneurysm had no pathological formations, there were no blood clots. The aneurysmically dilated left atrial appendage was adjacent to the trunk and left branch of the pulmonary artery, to the upper and lower left pulmonary veins, and to the left atrioventricular sulcus. The lower contour of the aneurysm reached the diaphragmatic surface of the pericardium. The left atrium was of normal size (Figure 1). Preoperative coronary angiography showed the absence of coronary artery stenosis.

---

## FINAL DIAGNOSIS

---

The final diagnosis of the presented case is left atrial appendage aneurysm.

---

## TREATMENT

---

The operative approach was *via* a median sternotomy with cardiopulmonary bypass. Along the lateral wall of the left atrium, in the projection of the left atrial appendage, there was a pulsating, spherical, and irregularly shaped aneurysm of 10 cm × 8 cm in size. It was not adjacent to the pericardium and involved the posterior wall, reaching the left pulmonary veins ostia.


The intraoperative transesophageal echocardiography showed that the aneurysmal cavity was divided by a short septum and had free communication with the left atrial cavity. Relative mitral insufficiency requiring correction was also revealed.

The mitral valve was accessed through the right contour of the left atrium. The aneurysm of the left atrium was opened from the outside, and its base was resected. The wall of the aneurysm was thin, flabby, and without any blood clots.


In the area of the circumflex artery, the left ventricle was significantly deformed and had an indentation of 3.0 cm × 2.0 cm in size. The lower edge of the resected aneurysm was located near the fibrous annulus of the mitral valve, in the area of anterolateral commissure (A1 and P1 segments). The flaps of the mitral valve and chords were not damaged. Mitral valve annuloplasty was performed by using a PROFILE 3D® 680R annuloplasty ring (Medtronic, United States). Considering that the defect, which formed after excision of the aneurysm, was close to the fibrous annulus and the circumflex artery, we decided to suture it from the inside. In the area of the fibrous annulus, the suture was fixed to the outer layer of the annuloplasty ring (Figure 2). Intraoperative transesophageal echocardiography showed grade 0-1 mitral regurgitation and insignificant mitral regurgitant volume.

Considering the anatomical location of the aneurysm and the absence of visualization of the circumflex artery, we performed coronary angiography. Signs of impaired blood flow in the left main coronary and circumflex artery and 60% stenosis were found. Bifurcation stenting was performed.


A histological examination of the resected left atrial appendage revealed a 1 mm thin wall of the left atrium with hypertrophied cardiomyocytes with dystrophic changes and vacuolated cytoplasm, and slight fibrosis of the endo- and epicardium (Figure 3).



**Figure 1 Magnetic resonance imaging.** A white arrow shows a sharply dilated cavity of the left atrial appendage.



**Figure 2 Aneurysmectomy.** A: The aneurysm of the left atrial appendage is opened; B: The aneurysm of the left atrial appendage is excised and sutured from the inside.



**Figure 3 Hematoxylin and eosin staining of the resected left atrial appendage aneurysm.** A: The wall of the aneurysm is thin, with slight fibrosis of the endo- and epicardium (scale bar 100  $\mu$ m, 100  $\times$  magnification); B: Hypertrophied cardiomyocytes with dystrophic changes and vacuolated cytoplasm (scale bar 100  $\mu$ m, 400  $\times$  magnification).

## OUTCOME AND FOLLOW-UP

The patient had an uneventful postoperative clinical course and was discharged from the hospital on the 10<sup>th</sup> d in a satisfactory condition. Sinus rhythm restored in the early postoperative period. Atrial flutter reappeared 2 mo after discharge from the hospital and required surgical intervention; so, radiofrequency catheter ablation of the cavotricuspid isthmus was performed. Transthoracic echocardiography performed 3 mo after discharge showed a left ventricular systolic pressure of 29 mmHg and an ejection fraction of 60%, grade 0-1 mitral regurgitation, and insignificant mitral regurgitant volume. The left atrium was 43 mm  $\times$  53 mm in size.

## DISCUSSION

In 1938, Semans and Taussig reported an isolated expansion of the cavity of the left atrium in a 5-year-old girl<sup>[2]</sup>. In 1962, Parmley described aneurysmal dilatation of the left atrium appendage in two children<sup>[3]</sup>. The first child of 11 years old had supraventricular cardiac arrhythmia and two episodes of systemic embolism and underwent resection of the left atrial appendage and removal of the atrial septal defect at the Mayo Clinic. The second, 7-year-old patient had an aneurysm of the left atrial appendage and a concomitant congenital anomaly of the left renal artery and left kidney causing arterial hypertension. In 1963, Williams published a report on an increase in the thin-walled appendage of the left atrium in a 27-year-old man<sup>[4]</sup>. An attempt to clamp the appendage during surgery was accompanied by cardiac arrhythmia; therefore, appendage resection had to be abandoned. However, due to embolic complications that occurred two years later, the removal of the left atrial appendage with blood clots was performed. To date, about 150 cases of this defect have been described in the literature<sup>[5,6]</sup>.

The defect has a congenital and acquired form. In the congenital form, the formation of the aneurysm occurs due to dysplasia of the scallop muscles and associated atrial muscle fibers<sup>[5]</sup>. Poor contractile function leads to a gradual expansion and extension of the left atrial appendage as a result of increased internal pressure<sup>[7]</sup>.

Foale *et al*<sup>[8]</sup> proposed the following diagnostic criteria for the congenital form: (1) Origin from an otherwise normal atrial chamber; (2) Well-defined communication with the atrial cavity; (3) Location within the pericardium; and (4) Distortion of the left ventricle by an aneurysm.

At the same time, Williams divided congenital aneurysms of the left atrium into extra- and intrapericardial types<sup>[4]</sup>. In the extrapericardial type, the left atrium or appendage is prolapsed through the pericardial defect and compressed in it, which leads to the aneurysmal expansion of the extrapericardial part. The intrapericardial type is due to the underdeveloped wall of the left atrium or its appendage<sup>[9]</sup>.

The process of development of acquired aneurysms is the result of damage to the mitral valve or other conditions leading to an increase in pressure in the left atrium<sup>[10]</sup>. At the same time, it is often difficult to differentiate the acquired and congenital forms, as in our case, since moderate changes of the mitral valve can be caused by the presence of the appendage aneurysm itself<sup>[11]</sup>.

Clinical signs of the disease occur in the third decade of life<sup>[1]</sup>. The most common manifestations are heart palpitations (43%), shortness of breath (22%), heart rhythm disturbances (15%), embolic disorders of cerebral circulation (11%), and chest pain and discomfort (7%)<sup>[12]</sup>.

Heart rhythm disturbances occur as a result of structural remodeling of the left atrial appendage, leading to electrical dissociation of cardiomyocytes and local disturbances in conduction that favor the emergence and maintenance of arrhythmia<sup>[13]</sup>. Left atrial appendage dilatation, atrial fibrillation, blood stasis, and thrombosis of the aneurysmal cavity are responsible for subsequent thromboembolic complications<sup>[5]</sup>.

Despite the large size of the aneurysm, our patient did not have any of the following symptoms associated with compression of surrounding structures: (1) Compression of the left ventricle by the aneurysm may cause an increase in filling pressure and diastolic dysfunction, leading to the appearance of shortness of breath and heart failure<sup>[13,14]</sup>; (2) Compression of the left anterior descending artery may manifest as angina pectoris<sup>[15]</sup>; (3) A characteristic sign of irritation of the left phrenic nerve is hiccups<sup>[6]</sup>; and (4) The aneurysm proximity to the respiratory tract may cause a dry, unproductive cough<sup>[16]</sup>.

Cases of left atrial appendage aneurysm rupture with a lethal outcome have been described<sup>[9]</sup>.

In most cases, the disease is detected by chance because of a non-specific clinical picture. A convexity at the contour of the left atrium determined on a chest X-ray requires differential diagnosis with a pericardial cyst and heart and mediastinal tumors<sup>[10,11]</sup>.

Echocardiography is considered the main method for diagnosing left atrial appendage aneurysm and the presence of blood clots in it<sup>[1]</sup>. While, due to the limited echo window, the transthoracic examination does not always make it possible to accurately establish the diagnosis, the transesophageal approach may provide a more detailed and clear visualization<sup>[10]</sup>.

Computed tomography and MRI are useful in unclear cases. MRI has a high resolution, which makes it optimal in assessing the surrounding structures and cardiac abnormalities. However, MRI has certain disadvantages, such as the need for regular

heart rhythm and patient's exposure to nephrotoxic contrast agents. Cardiac computed tomography makes it possible to evaluate the anatomy of the coronary arteries, if there is a suspicion of their compression, but cannot provide functional data as accurately as echocardiography or MRI<sup>[10]</sup>.

Surgical treatment is indicated for all patients to prevent the occurrence of atrial fibrillation, systemic embolism, or myocardial dysfunction<sup>[17,18]</sup>. We performed a left atrial appendage aneurysm resection *via* median sternotomy with cardiopulmonary bypass, which is considered the operation of choice, especially in patients with thrombosis of the aneurysmal cavity and the need for concomitant interventions<sup>[19]</sup>.

Aneurysmectomy through left-sided lateral thoracotomy should be considered if isolated lesions of the left atrial appendage are present without significant violations of the heart anatomy<sup>[20]</sup>. Endoscopic surgery is most suitable for patients with minor aneurysms in the absence of thrombosis<sup>[21]</sup>.

The issue of performing a concomitant intervention to restore sinus rhythm remains debatable. Following the most recent 2017 guidelines from the Society of Thoracic Surgeons, surgical ablation for atrial fibrillation is recommended at the time of concomitant mitral operations to restore sinus rhythm (Class I, Level A)<sup>[22]</sup>. However, removal of the aneurysm usually results in resolution of atrial arrhythmias with no need for additional surgery<sup>[23,24]</sup>. Moreover, concomitant atrial fibrillation surgery increases the risk for requiring a permanent pacemaker<sup>[25]</sup>. We decided to abandon concomitant surgery for atrial fibrillation to limit the overall risks of the operation. Also, there was a possibility that atrial flutter might disappear after resection of the aneurysm. Unfortunately, arrhythmia disappeared only temporarily and required radiofrequency catheter ablation 2 mo after discharge from the hospital.

### Limitations

This is a case report and the conclusions are subject to the inherent bias associated with the retrospective design. Also, this study design does not allow us to establish a cause-effect relationship between atrial appendage aneurysm and mitral regurgitation, as well as to differentiate between congenital and acquired forms of the aneurysm.

---

## CONCLUSION

Left atrial appendage aneurysm is a rare heart defect. Given the nonspecific clinical picture, the disease is diagnosed either by chance or during the examination of a patient when complications arise. Echocardiography helps to establish the diagnosis in most of the cases. However, MRI is necessary for a more detailed assessment of changes in the heart. Surgical treatment is indicated for all patients to prevent fatal disorders. In the case when the circumflex artery is not visualized, coronary angiography may help to avoid adverse events. Antiarrhythmic surgery can decrease the risk of thromboembolic complications and may be performed either at the time of concomitant left atrial appendage resection or as a stand-alone surgical procedure, depending on the clinical situation.

---

## REFERENCES

- 1 **Chen Y**, Mou Y, Jiang LJ, Hu SJ. Congenital giant left atrial appendage aneurysm: a case report. *J Cardiothorac Surg* 2017; **12**: 15 [PMID: 28302138 DOI: 10.1186/s13019-017-0576-6]
- 2 **Semans JH**, Taussig HB. Congenital "aneurysmal" dilatation of the left auricle. *Bull Johns-Hopkins Hosp* 1938; **63**: 404
- 3 **Parmley LF**. Congenital atriomegaly. *Circulation* 1962; **25**: 553-558 [PMID: 14483987 DOI: 10.1161/01.cir.25.3.553]
- 4 **Williams WG**. Dilatation of the Left Atrial Appendage. *Br Heart J* 1963; **25**: 637-643 [PMID: 14063011 DOI: 10.1136/hrt.25.5.637]
- 5 **Aryal MR**, Hakim FA, Ghimire S, Ghimire S, Giri S, Pandit A, Bhandari Y, Bhandari N, Pathak R, Karmacharya P, Pradhan R. Left atrial appendage aneurysm: a systematic review of 82 cases. *Echocardiography* 2014; **31**: 1312-1318 [PMID: 24976376 DOI: 10.1111/echo.12667]
- 6 **Asfalou I**, Boumaaz M, Raissouni M, Sabry M, Benyass A, Zbir EM. Huge left atrial appendage aneurysm revealed by chronic hiccups. *J Saudi Heart Assoc* 2017; **29**: 293-296 [PMID: 28983173 DOI: 10.1016/j.jsha.2017.03.009]
- 7 **Wang HQ**, Zhang Z, Yang H, Wu S, Fu YH, Song ZM, Wan F. A Huge Congenital Left Atrial Appendage Aneurysm. *Chin Med J (Engl)* 2017; **130**: 3011-3012 [PMID: 29237940 DOI: 10.4103/0366-6999.220301]
- 8 **Foale RA**, Gibson TC, Guyer DE, Gillam L, King ME, Weyman AE. Congenital aneurysms of the left atrium: recognition by cross-sectional echocardiography. *Circulation* 1982; **66**: 1065-1069
- 9 **Chowdhury UK**, Seth S, Govindappa R, Jagia P, Malhotra P. Congenital left atrial appendage aneurysm: a

- case report and brief review of literature. *Heart Lung Circ* 2009; **18**: 412-416 [PMID: 19119072 DOI: 10.1016/j.hlc.2008.10.015]
- 10 **Wang B**, Li H, Zhang L, He L, Zhang J, Liu C, Wang J, Lv Q, Shang X, Liu J, Xie M. Congenital left atrial appendage aneurysm: A rare case report and literature review. *Medicine (Baltimore)* 2018; **97**: e9344 [PMID: 29480827 DOI: 10.1097/MD.0000000000009344]
  - 11 **Valentino MA**, Al Danaf J, Morris R, Tecce MA. Giant left atrial appendage aneurysm: A case of mistaken identity. *J Cardiol Cases* 2017; **15**: 129-131 [PMID: 30279759 DOI: 10.1016/j.jccase.2016.12.010]
  - 12 **Bamous M**, Aithoussa M, Abetti A, Boulahya A. Congenital left atrial appendage aneurysm: Atypical presentation. *Ann Pediatr Cardiol* 2017; **10**: 293-294 [PMID: 28928618 DOI: 10.4103/apc.APC\_4\_17]
  - 13 **Li M**, Wang Y, Liu S, Yang J, Ma C. Giant left atrial appendage aneurysm compressing the left ventricular wall diagnosed by multiple imaging technology. *Cardiol J* 2019; **26**: 416-417 [PMID: 31452186 DOI: 10.5603/CJ.2019.0079]
  - 14 **Harland DR**, Suma V, Muthukumar L, Port SC, Werner PH, Tajik AJ. Giant Congenital Left Atrial Appendage Aneurysm Presenting With Recurrent Supraventricular Tachycardia and Chest Pain. *CASE (Phila)* 2019; **3**: 129-132 [PMID: 31286094 DOI: 10.1016/j.case.2019.01.003]
  - 15 **Wagdy K**, Samaan A, Romeih S, Simry W, Afifi A, Hassan M. Giant left atrial appendage aneurysm compressing the left anterior descending coronary artery. *Echocardiography* 2016; **33**: 1790-1792 [PMID: 27539326 DOI: 10.1111/echo.13296]
  - 16 **Toufan M**, Pourafkari L, Afrasiabi A, Sohrabi M, Nader ND. Left atrial appendage aneurysm presenting with chronic cough. *Neth Heart J* 2017; **25**: 526-527 [PMID: 28730519 DOI: 10.1007/s12471-017-1021-0]
  - 17 **Teng P**, Ni Y, Sun Q, Zhao H. Giant Left Atrial Appendage Aneurysm: Surgical Treatment to Prevent Potential Complications. *Heart Surg Forum* 2018; **21**: E464-E465 [PMID: 30604668 DOI: 10.1532/hcf.2129]
  - 18 **Lukin OP**, Belov DV, Milievskaia EB. [Organization of the cardiosurgical medical care in the Ural Federal District]. *Grudnaya i serd-sosud hir* 2018; **60**: 281-286 [DOI: 10.24022/0236-2791-2018-60-4-281-286]
  - 19 **Victor S**, Nayak VM. Aneurysm of the left atrial appendage. *Tex Heart Inst J* 2001; **28**: 111-118 [PMID: 11453121]
  - 20 **Clark JB**, Ting JG, Polinsky RJ Jr, Wolfe LT. Resection of a Giant Left Atrial Appendage Aneurysm via Limited Thoracotomy. *World J Pediatr Congenit Heart Surg* 2014; **5**: 475-477 [PMID: 24958057 DOI: 10.1177/2150135114524602]
  - 21 **Ruttkey T**, Scheid M, Götte J, Doll N. Endoscopic Resection of a Giant Left Atrial Appendage. *Innovations (Phila)* 2015; **10**: 282-284 [PMID: 26355691 DOI: 10.1097/IML.0000000000000172]
  - 22 **Badhwar V**, Rankin JS, Damiano RJ Jr, Gillinov AM, Bakaeen FG, Edgerton JR, Philpott JM, McCarthy PM, Bolling SF, Roberts HG, Thourani VH, Suri RM, Shemin RJ, Firestone S, Ad N. The Society of Thoracic Surgeons 2017 Clinical Practice Guidelines for the Surgical Treatment of Atrial Fibrillation. *Ann Thorac Surg* 2017; **103**: 329-341 [PMID: 28007240 DOI: 10.1016/j.athoracsur.2016.10.076]
  - 23 **Burke RP**, Mark JB, Collins JJ Jr, Cohn LH. Improved surgical approach to left atrial appendage aneurysm. *J Card Surg* 1992; **7**: 104-107 [PMID: 1606362 DOI: 10.1111/j.1540-8191.1992.tb00786.x]
  - 24 **Vagefi PA**, Choudhry M, Hilgenberg AD. Excision of an aneurysm of the left atrial appendage. *J Thorac Cardiovasc Surg* 2007; **133**: 822-823 [PMID: 17320601 DOI: 10.1016/j.jtcvs.2006.11.025]
  - 25 **Huffman MD**, Karmali KN, Berendsen MA, Andrei AC, Kruse J, McCarthy PM, Malaisrie SC. Concomitant atrial fibrillation surgery for people undergoing cardiac surgery. *Cochrane Database Syst Rev* 2016; CD011814 [PMID: 27551927 DOI: 10.1002/14651858.CD011814.pub2]



Published by **Baishideng Publishing Group Inc**  
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA  
**Telephone:** +1-925-3991568  
**E-mail:** [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)  
**Help Desk:** <https://www.f6publishing.com/helpdesk>  
<https://www.wjgnet.com>

